

OCHA-PP-125
NDA-FP-51
October 1998

D0-brane description of the charged black hole

YURIKO KATO¹
SHIN'ICHI NOJIRI^{♣2} and AKIO SUGAMOTO³

*Department of Physics, Ochanomizu University
Otsuka, Bunkyo-ku Tokyo 112, JAPAN*

[♣] *Department of Mathematics and Physics
National Defence Academy, Hashirimizu Yokosuka 239, JAPAN*

ABSTRACT

The charged black hole is considered from the viewpoint of D0-brane in the Matrix theory. It can be obtained from the Kaluza-Klein mechanism by boosting the Schwarzschild black hole in a circle, which is the compactified one dimensional space. Especially, how the extremal limit is realized by the Boltzmann gas of D0-brane, has been shown. In the course of our discussion, the Virial theorem for the statistical average plays an important role.

PACS:

¹e-mail : g9740507@edu.cc.ocha.ac.jp

² e-mail : nojiri@cc.nda.ac.jp

³ e-mail : sugamoto@phys.ocha.ac.jp

It is known that the several properties of the Schwarzschild black holes can be well described by the Boltzmann gas of the D0-brane [1, 2], the parton of the Matrix theory [3]. Recently Ohta and Zhou [4] have calculated the partition function of the Matrix theory in the D0-brane background. They have derived the mass, entropy, temperature etc. of this system and shown that the D0-brane background can be interpreted as 11D Schwarzschild black hole.

Since D0-branes interact with each other very weakly, the most of the properties of the D0-brane background can be derived from the description of the free Boltzmann gas. The information of the interaction can be included by using the Virial theorem.

In this letter, we treat the charged black hole in the Matrix theory, which can be obtained from the Kaluza-Klein mechanism by boosting the Schwarzschild black hole in a circle, which is the compactified one dimensional space. Especially, we show how the extremal limit is described by the Boltzmann gas of D0-brane. In [4], the oscillation modes in the Matrix theory is integrated out by using the Euclidean path integral method. In this letter, we give the more statistical method for the treatment of the zero modes, which reproduces the same results of the Schwarzschild black hole in [4] derived from the consistency, and gives the new results on the charged black hole.

We start with the free part of the M-theory effective Lagrangian in [4], which is given by integrating oscillating modes in a given background $\bar{X}^i$:

$$\mathcal{L}^{\text{free}} = \text{Tr} \frac{1}{2R} (\dot{\bar{X}}^i)^2, \quad (1)$$

where R is the radius of the light-like compactification from 11D to 10D. Then the canonical momentum P^i conjugate to $\bar{X}^i$ reads

$$P^i = \frac{1}{R} \dot{\bar{X}}^i. \quad (2)$$

As a background $\bar{X}^i$, we consider the following one in which N D0-branes are moving freely :

$$\bar{X}^i = \text{diag.}(v_1^i t + b_1^i, v_2^i t + b_2^i, \dots, v_N^i t + b_N^i). \quad (3)$$

If the black hole is a statistical system in which D0-branes are moving inside the sphere of the radius R_s at temperature β^{-1} in the light-cone frame, we

can define the partition function of the black hole as follows:⁴

$$\begin{aligned}
Z(\beta, N) &= \int d\bar{X}^i e^{-\int_{-\frac{\beta}{2}}^{\frac{\beta}{2}} \mathcal{L}^{\text{free}}} \\
&\sim \frac{1}{R^{9N}} \int_{-\infty}^{\infty} \prod dv_n^i \int_{-\frac{R_s}{2}}^{\frac{R_s}{2}} \prod db_n^i e^{-\frac{\beta}{2R} (v_n^i)^2} \\
&= \left(\frac{2\pi R_s^2}{\beta R} \right)^{\frac{9N}{2}}.
\end{aligned} \tag{4}$$

Note that, from (2), the phase space volume for a pair of n -th coordinate and its conjugate momentum of the background D0-brane is given by (we choose $\hbar = 1$)

$$dx_n^i dp_n^i = \frac{dv_n^i db_n^i}{R}. \tag{5}$$

By using (4), we find

$$\begin{aligned}
\langle v_n^2 \rangle &= -\frac{2R}{N} \frac{\partial}{\partial \beta} \ln Z(\beta, N) \\
&= \frac{9R}{\beta}.
\end{aligned} \tag{6}$$

Since β is the Euclidean time interval, the typical magnitude of the velocity of D0-brane moving within the black hole having the radius R_s is given by

$$\langle v_n^2 \rangle \sim \frac{R_s^2}{\beta^2}. \tag{7}$$

By comparing (6) and (7), we find

$$\beta \sim \frac{R_s^2}{R}, \tag{8}$$

which corresponds to Eq.(2.9) in ref.[4]. By using (4), we find the Helmholtz free energy $F(\beta, N)$

$$\begin{aligned}
F(\beta, N) &= -\frac{1}{\beta} \ln Z(\beta, N) \\
&= -\frac{9N}{2\beta} \ln \left(\frac{2\pi R_s^2}{\beta R} \right),
\end{aligned} \tag{9}$$

⁴The explicit integrations for the zero modes v_n^1 and b_n^i have been not given in [4].

and the entropy S

$$\begin{aligned} S(\beta, N) &= \beta^2 \frac{\partial F}{\partial \beta} \\ &= \frac{9N}{2} \left\{ \ln \left(\frac{2\pi R_s^2}{\beta R} \right) + 1 \right\} . \end{aligned} \quad (10)$$

From (8), we obtain

$$S(\beta, N) \sim N \quad (11)$$

which corresponds to (3.11) in [4]. The expectation value of the energy E is also given by

$$\begin{aligned} E &= -\frac{\partial}{\partial \beta} \ln Z(\beta, N) \\ &= \frac{9N}{2\beta} . \end{aligned} \quad (12)$$

We now use the expression (3.1) in [4] for E

$$\begin{aligned} E &= \sum_{n=1}^N \frac{v_n^2}{2R} - \frac{15G_{11}}{16R^3} \sum_{n < m}^N \frac{v_{nm}^4}{\left(b_{nm}^2 + \frac{v_{nm}^2 \beta^2}{4} \right)^{\frac{7}{2}}} \\ v_{nm}^i &\equiv v_n^i - v_m^i \\ b_{nm}^i &\equiv b_n^i - b_m^i . \end{aligned} \quad (13)$$

This is the effective energy for a given background configuration, including the fluctuations around it, but the statistical averaging over the background configurations is further required. Therefore we apply the Virial theorem to (13) by regarding E in (13) as the Hamiltonian and assume

$$v_n^2 \sim v_{nm}^2 \sim \frac{R_s^2}{\beta^2} \sim \frac{R}{\beta} , \quad b_{nm}^2 \sim R_s^2 \quad (14)$$

as in (7). Then we find

$$R_s \sim (G_{11}N)^{\frac{1}{9}} , \quad (15)$$

The derivation of which will be clarified in the following⁵. By using (8), (12) and (15), we find also

$$E \sim \frac{\left(G_{11}^{-\frac{1}{9}} N^{\frac{8}{9}}\right)^2 R}{N} . \quad (16)$$

The relations (15) and (16) correspond to (3.9) in [4].

Usually the Virial theorem is given by the average with respect to time. In the language of the statistical average, the theorem can be proved in the following way: Here we only consider the case in that Hamiltonian H is given by a pair of one coordinate q and its conjugate momentum p :

$$H = H(q, p) , \quad (17)$$

since the generalization to the case with several coordinates and their conjugate momenta is straightfoward. We can observe that the expectation value of $p \frac{\partial H}{\partial p} - q \frac{\partial H}{\partial q}$ vanishes, since it is the total derivative in the phase space

$$\begin{aligned} & \left\langle p \frac{\partial H}{\partial p} - q \frac{\partial H}{\partial q} \right\rangle \\ &= \frac{1}{Z(\beta)} \int dq dp \left(p \frac{\partial H}{\partial p} - q \frac{\partial H}{\partial q} \right) e^{-\beta H(q,p)} \\ &= \frac{1}{Z(\beta)} \int dq dp \left\{ \frac{\partial}{\partial p} (p e^{-\beta H(q,p)}) - \frac{\partial}{\partial q} (q e^{-\beta H(q,p)}) \right\} \\ &= 0 , \end{aligned} \quad (18)$$

where

$$Z(\beta) \equiv \int dq dp e^{-\beta H(q,p)} . \quad (19)$$

The first term $T = p \frac{\partial H}{\partial p}$ corresponds to the kinetic energy and the second term $V = q \frac{\partial H}{\partial q}$ to the so-called “virial”, so that we find the Virial theorem as

$$\langle T \rangle = \langle V \rangle . \quad (20)$$

By applying the above theorem to (13) and using (14), we find (15).

⁵Essentially we assume that the expectation values of the first and the second terms are the same magnitude with each other.

Now we extend the above analysis to the case of charged black hole. The charged black hole can be obtained by the Kaluza-Klein mechanism [5]. We impose a periodic boundary condition on one dimension in 11 dimensional Schwarzschild black hole spacetime and compactify it to a circle. After that, we boost the black hole solution in that one dimensional direction. Then the total momentum in the compactified space can be regarded as the total charge of the black hole.

Let X^1 be the coordinate in the compactified space. We now project the partition function in (4) to the sector in which total momentum conjugate to X^1 is $\frac{V}{R}$, by inserting a delta function. By using the following expression of a delta function

$$\delta\left(\sum_{n=1}^N v_n^1 - V\right) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ik(\sum_{n=1}^N v_n^1 - V)}, \quad (21)$$

we obtain

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\infty}^{\infty} \prod_{n=1}^N dv_n^1 \int_{-\infty}^{\infty} dk e^{-\frac{\beta}{2R} \sum_{n=1}^N (v_n^1)^2 + ik(\sum_{n=1}^N v_n^1 - V)} \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \prod_{n=1}^N dv_n^1 \int_{-\infty}^{\infty} dk e^{-\frac{\beta}{2R} \sum_{n=1}^N (v_n^1 - \frac{ikR}{\beta})^2 - \frac{NR}{2\beta} (k + \frac{i\beta V}{NR})^2 - \frac{\beta V^2}{2NR}} \\ &= \left(\frac{\beta}{2\pi NR}\right)^{\frac{1}{2}} e^{-\frac{\beta V^2}{2NR}} \int_{-\infty}^{\infty} \prod_{n=1}^N dv_n^1 e^{-\frac{\beta}{2R} \sum_{n=1}^N (v_n^1)^2} \end{aligned} \quad (22)$$

and we find that the projected partition function is given by

$$Z(\beta, N, V) = \left(\frac{2\pi R_s^2}{\beta R}\right)^{\frac{9N}{2}} \left(\frac{\beta}{2\pi NR}\right)^{\frac{1}{2}} e^{-\frac{\beta V^2}{2NR}}. \quad (23)$$

In a similar way to derive (6), from the partition function in (23), we obtain

$$\langle v_n^2 \rangle = \left(9R - \frac{R}{N}\right) \frac{1}{\beta} + \frac{V^2}{N^2}. \quad (24)$$

By assuming $\langle v_n^2 \rangle \sim \frac{R_s^2}{\beta^2}$ as in (7), we can solve Eq.(24) with respect to β and obtain

$$\frac{1}{\beta} = \frac{9R \pm \sqrt{(9R)^2 + \frac{4R_s^2 V^2}{N^2}}}{2R_s^2}. \quad (25)$$

Here we also assume $N \gg 1$. In (25), the negative sign cannot be physically accepted since it gives the negative temperature. The expression of the entropy S in (10) does not be changed. The expectation value of the energy is given by

$$E = \frac{9N - 1}{2\beta} + \frac{V^2}{2NR} . \quad (26)$$

Since E is related with the black hole mass M [1] as follows

$$E = \frac{M^2 R}{2N} , \quad (27)$$

we find

$$M^2 = \frac{(9N - 1)N}{R\beta} + \frac{V^2}{R^2} . \quad (28)$$

Furthermore by using the relation between the light-cone temperature β^{-1} and the rest frame temperature T [4],

$$\frac{1}{\beta} = \frac{MR}{2N} T , \quad (29)$$

we find

$$T = \frac{2 \left(M^2 - \frac{V^2}{R^2} \right)}{(9N - 1)M} . \quad (30)$$

Remarkably Eq.(30) tells that the temperature vanishes when

$$M = Q \equiv \left| \frac{V}{R} \right| . \quad (31)$$

Therefore Eq.(31) is nothing but the condition of the extremal limit of the black hole.

In summary, we have derived the mass, temperature etc. of the charged black hole by using the D0-brane description in the Matrix theory. Especially, we have found how the extremal limit is described.

References

- [1] T. Banks, W. Fischler, I. Klebanov, L. Susskind, *Phys.Rev.Lett.* 80 (1998) 226, *J.High Energy Phys.* **01** (1998) 008.

- [2] M. Li, *J.High Energy Phys.* **01** (1998) 009.
- [3] T. Banks, W. Fischler, S.H. Shenker, L. Susskind, *Phys.Rev* **D55** (1997) 5112.
- [4] N. Ohta, J.-G. Zhou, *Nucl.Phys.* **B522** (1998) 125.
- [5] J.H. Horne, G.T. Horowitz, R. Steif, *Phys.Rev.Lett.* **68** (1992) 568.